

MONODROMY WEIGHT FILTRATION IS INDEPENDENT OF l

TOMOHIKE TERASOMA

§1 WEIGHT MONODROMY FILTRATION FOR NILPOTENT MONODROMY

In this section, we recall several fundamental properties of monodromy weight filtration for nilpotent monodromy and prove some approximation theory. Let K_0 a local field of characteristic p with a finite residue field $\kappa = \mathbf{F}_q$. The integer ring of K_0 is denoted by R_0 . Let $\mathbf{F}$ be an algebraic closure of $\mathbf{F}_q$ and $R = R_0 \otimes_{\mathbf{F}_q} \mathbf{F}$. The fraction field of R is denoted by K and we use the notations

$$\begin{aligned} \operatorname{Spec}(R_0) &= S_0, \operatorname{Spec}(K_0) = \eta_0, \operatorname{Spec}(\kappa_0) = s_0, \\ \operatorname{Spec}(R) &= S, \operatorname{Spec}(K) = \eta, \operatorname{Spec}(\kappa) = s. \end{aligned}$$

The algebraic closure of K is denoted by $\bar{K}$ and $\operatorname{Spec}(\bar{K})$ is denoted by $\bar{\eta}$. Let $\mathcal{X}_0$ be a scheme over S_0 whose generic geometric fiber $X_{\bar{\eta}} = \mathcal{X}_0 \times_{S_0} \bar{\eta}$ is smooth. Denote by $\mathcal{X}$ the fiber product $\mathcal{X}_0 \times_{\mathbf{F}_q} \mathbf{F}$. Then by a theorem of Grothendieck, the action of the Galois group $I = \operatorname{Gal}(\bar{K}/K)$ on $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$ is quasi unipotent. Let J be an open compact subgroup of I whose action on $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$ is unipotent. The action of J factors through the maximal tame quotient J^t of J . Let U a topological generator of $J^t \simeq \hat{\mathbf{Z}}'(1)$ and N be the logarithm of the action of U on $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$. In [D], Deligne introduced an increasing filtration $W_\bullet$ with the following properties:

- (1) $N(W_k H^i(X_{\bar{\eta}}, \mathbf{Q}_l)) \subset W_{k-2} H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$
- (2) The induced map $N : Gr_k^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l)) \rightarrow Gr_{k-2}^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l))$ induces an isomorphism;

$$N^k : Gr_k^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l)) \xrightarrow{\cong} Gr_{-k}^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l)).$$

We define the primitive part $P_k(H^i(X_{\bar{\eta}}, \mathbf{Q}_l))$ of $Gr_k^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l))$ ($k \geq 0$) by the kernel of N^{k+1} . Let G be an open compact subgroup of $\operatorname{Gal}(\bar{K}/K_0)$ such that $J = G \cap \operatorname{Gal}(\bar{K}_0/K)$ acts $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$ nilpotently. Let L_0 be the corresponding extension of K_0 and $\mathbf{F}_{q'}$ be the residue field. Then we have the exact sequence

$$1 \rightarrow J \rightarrow G \rightarrow \operatorname{Gal}(\mathbf{F}/\mathbf{F}_{q'}) \rightarrow 1.$$

Since the action of J on the associated graded vector space $Gr_k^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l))$ is trivial, $\operatorname{Gal}(\mathbf{F}/\mathbf{F}_{q'})$ acts on this vector space.

Before studying the action of $\operatorname{Gal}(\mathbf{F}/\mathbf{F}_{q'})$ on $Gr_k^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l))$, we try to approximate the local situation by the global situation. Let $g_0 : Y_0 \rightarrow C_0$ be a projective generic geometrically smooth morphism of relative dimension n , where C_0 is a projective curve over $\mathbf{F}_q$. Let p be an $\mathbf{F}_q$ valued point of C_0 .

We denote by O_p and K_p by the completion of the structure sheaf of C_0 at the point p and its quotient field respectively. Let $\bar{K}_p$ be the algebraic closure of K_p , and $\bar{\xi} = \operatorname{Spec}(\bar{K}_p)$. We denote by $Y_{\bar{\xi}}$ the base extension of Y_0 to $\bar{\xi}$.

Lemma 1.1. *Let $f_0 : \mathcal{X}_0 \rightarrow S_0$ be as above and $l_1, \dots, l_s$ be finite set of primes different from p . There exist a projective smooth curve C_0 over $\mathbf{F}_q$, and a projective geometric generically smooth morphism $Y_0 \rightarrow C_0$ of relative dimension n with the following property:*

(1) *There exists an isomorphism*

$$(1.1) \quad H^i(Y_{\bar{\xi}}, \mathbf{Q}_{l_i}) \simeq H^i(X_{\bar{\eta}}, \mathbf{Q}_{l_i})$$

for all $i = 1, \dots, s$.

(2) *There exists an isomorphism*

$$\begin{array}{ccc} \mathrm{Gal}(\bar{K}_0/K_0)/P_{K_0,i} & \longrightarrow & \mathrm{Gal}(\bar{\mathbf{F}}_q/\mathbf{F}_q) \\ \simeq \downarrow & & \downarrow = \\ \mathrm{Gal}(\bar{K}_p/K_p)/P_{K_p,i} & \longrightarrow & \mathrm{Gal}(\bar{\mathbf{F}}_q/\mathbf{F}_q), \end{array}$$

which is compatible with the isomorphism of (1.1), where $P_{K_0,i}$ and $P_{K_p,i}$ is the kernel of

$$\mathrm{Gal}(\bar{K}_0/K) \rightarrow \mathrm{Aut}(H^i(X_{\bar{\eta}}, \mathbf{Q}_{l_i}))$$

and

$$\mathrm{Gal}(\bar{K}_p/K_p \otimes_{\mathbf{F}_q} \mathbf{F}) \rightarrow \mathrm{Aut}(H^i(Y_{\bar{\xi}}, \mathbf{Q}_{l_i}))$$

respectively.

Proof. Let us write $R_0 = \mathbf{F}_q[[t]]$. By fixing a projective model of $\mathcal{X}_0$ on S_0 , we can take a finitely generated sub ring $\mathbf{F}_q[t, x_1, \dots, x_m]$ of $\mathbf{F}_q[[t]]$ which contains all the coefficients of $\mathcal{X}_0$. The element corresponding to x_i is denoted by $\xi_i(t) \in \mathbf{F}_q[[t]]$. Let $M_0 = \mathrm{Spec}(\mathbf{F}_q[t, x_1, \dots, x_m])$. By the definition of M_0 there exists a projective variety $\mathcal{X}_{M_0} \rightarrow M_0$ of relative dimension n such that the base change of $\mathcal{X}_{M_0}$ by the map $S_0 \rightarrow M_0$ is $\mathcal{X}_0$. Let $M = M_0 \otimes_{\mathbf{F}_q} \mathbf{F}$, $\mathcal{X}_M = \mathcal{X}_{M_0} \otimes_{\mathbf{F}_q} \mathbf{F}$, and $f_M : \mathcal{X}_M \rightarrow M$. Since it is generically smooth, there exists an open set U_0 of M_0 such that $R^i(f_M)_*$ is a smooth sheaf on $U = U_0 \otimes_{\mathbf{F}_q} \mathbf{F}$. The complement of U_0 in M_0 is denoted by D_0 . By taking a finite covering of M_0 and alterations [J], Theorem 7.3, we have a proper dominant morphism $\phi : N_0 \rightarrow M_0$ and an action of a group G on N_0 over M_0 with the following property:

- (1) Let $Q(N_0)$ and $Q(M_0)$ be the rational function field of N_0 and M_0 . The field extension $Q(N_0)^G$ of $Q(M_0)$ is purely inseparable.
- (2) The variety N_0 is smooth and $E_0 = \phi^{-1}(D_0)$ is a G -strict normal crossing divisor.
- (3) Let $\mathbf{F}_{q'}$ be the constant field of N_0 , $N = N_0 \otimes_{\mathbf{F}_{q'}} \mathbf{F}$, $E = E_0 \otimes_{\mathbf{F}_{q'}} \mathbf{F}$, $f_N : \mathcal{X}_N \rightarrow N$ be the base change of f_M by the morphism $N \rightarrow M$. Let $\bar{\delta}$ be a geometric point of $N - E$. Then the image of the geometric monodromy action of $\pi_1(N - E, \bar{\delta})$ on the geometric fiber of $R^i(f_N)_* \mathbf{Q}_{l_i}$ at $\bar{\delta}$ is a pro- l_i group.

Since the field $Q(N_0)$ is a finite extension of $Q(M_0)$, by choosing an embedding of $Q(N_0)$ into the algebraic closure of K , the composite field $L = K \cdot Q(N_0)$ is

a finite extension of K_0 . Let T_0 be the trait corresponding to the extension L_0 . Then, by using valuative criterion of properness, we have the following diagram.

$$(1.2) \quad \begin{array}{ccc} T_0 & \longrightarrow & N_0 \\ \downarrow & & \downarrow \\ S_0 & \longrightarrow & M_0 \end{array}$$

Let t_0 be the image of the closed point of T_0 in N_0 . Let G_0 be the subgroup of G which preserves the image of T_0 . By blowing up t_0 , we get the similar diagram as (1.2). Repeating the process, we may assume that the image of the trait T_0 intersects only one component $E_{0,i}$ of E_0 transversally. Let m be the maximal ideal corresponding to t_0 and $\hat{N}_{0,m}$ be the completion of N_0 at m and $\hat{N}_m = \hat{N}_{0,m} \times_{\mathbf{F}_{q'}} \mathbf{F}$. Since the image of T_0 meets $E_{0,i}$ transversally, the natural homomorphism

$$\pi_1(T - \{t\})^{(l_i)} \simeq \mathbf{Z}_{l_i}(1) \rightarrow \pi_1(\hat{N}_m - E_i)^{(l_i)} \simeq \mathbf{Z}_{l_i}(1)$$

is an isomorphism, where $T = T_0 \times_{\mathbf{F}_{q'}} \mathbf{F}$ and $E_i = E_{0,i} \times_{\mathbf{F}_{q'}} \mathbf{F}$. Here we denote by $\pi_1(T - \{t\})^{(l)}$ the maximal pro- l quotient of the fundamental group of $T - \{t\}$. Now we take a curve D_0 in N_0 passing through t_0 meeting $E_{0,i}$ transversally and equivariant under the action of G_0 . Let $Z_0 = (f_{N_0})^{-1}(C_0)$ and take a quotient $Y_0 = Z_0/G_0$ and $C_0 = D_0/G_0$ of Z_0 and D_0 under the action of G_0 . Let p_0 be the image of t_0 under the quotient map. Then the induced morphisms $Y_0 \rightarrow C_0$ and p_0 satisfies the required properties.

As an application of this approximation lemma, we have the following Deligne's theorem for varieties on local fields.

Lemma 1.2 (see also [D]). *The action of $\text{Gal}(\mathbf{F}/\mathbf{F}_{q'})$ on $Gr_k^W(H^i(X_{\bar{\eta}}, \mathbf{Q}_l))$ is of pure weight $i + k$.*

Definition. The filtration W introduced here is called the weight monodromy filtration of $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$.

§2 INDEPENDENCE OF l IN THE GLOBAL SITUATION

Let C_0 be a proper smooth curve over $\mathbf{F}_q$, $f_0 : X_0 \rightarrow C_0$ be a projective flat relative n -dimensional morphism with generic geometrically smooth fiber. Let $X = X_0 \otimes_{\mathbf{F}_q} \mathbf{F}$, $C = C_0 \otimes_{\mathbf{F}_q} \mathbf{F}$, and K_0 and K be the function field of C_0 and C respectively. Let s_0 be a $\mathbf{F}_q$ -valued point of C_0 and s be the spectrum of the algebraic closure of the residue field of s_0 . For a rational function $g \in K$ and a character $\chi : \mu_d(\mathbf{F}) \rightarrow \mathbf{Q}_l^\times$, the associate Kummer sheaf on C is denoted by $\text{Kum}(g, \chi)$. The twist of χ by an element of Galois group $\sigma \in \text{Gal}(\mathbf{F}_q(\mu_d)/\mathbf{F}_q)$ is denoted by χ^σ . Let $\mathcal{F}_0$ be an etale sheaf on C_0 with a finite geometric monodromy. The restriction of $\mathcal{F}_0$ to C is denoted by $\mathcal{F}$.

Lemma 2.1. *There exists an element $g \in K_0$, a positive integer d , and an injective character $\chi : \mu_d \rightarrow \mathbf{Q}_l^\times$ such that*

- (1) *the support $\text{Supp}(g)$ of the divisor (g) in C is non-empty and the finite set $\text{Supp}(f) \cup \{s_0\}$ contains all the points in C where the morphism f or the sheaf $\mathcal{F}$ are not smooth.*

- (2) $f(s_0) = 1$,
- (3) the order of the class g in $K^\times/K^{d\times}$ is d , and
- (4) for all $x \in \text{Supp}(g)$ and $\sigma \in \text{Gal}(\mathbf{F}_{q'}/\mathbf{F}_q)$, the semi-simplification of the local monodromy group on $R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi^\sigma) \otimes \mathcal{F}$ at x has no fixed part.

Proof. Fix one closed point x_1 in C_0 different from s_0 . By Riemann-Roch theorem, we can choose a rational function g on C_0

- (1) whose order at x_1 is 1,
- (2) if f or $\mathcal{F}$ are not smooth at a point x different from s_0 , then $\text{ord}_x(g) \neq 0$, and
- (3) regular at s_0 and $g(s_0) = 1$.

Choose a sufficiently big d and χ such that $\text{Kum}(g, \chi)$ has the property (4). Then we get the required g , d , and χ .

Remark 2.2. The condition (1) and (4) of Lemma 2.1 implies

$$H_c^0(U, R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi^\sigma) \otimes \mathcal{F}) = H_c^2(U, R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi^\sigma) \otimes \mathcal{F}) = 0$$

for all $\sigma \in \text{Gal}(\mathbf{F}_{q'}/\mathbf{F}_q)$.

Now we introduce a covering $\tilde{C}_0, \tilde{X}_0$ of C_0 and X_0 . Let $\mathbf{F}_{q'} = \mathbf{F}_q(\mu_d)$ and $C_1 = C_0 \otimes_{\mathbf{F}_q} \mathbf{F}_{q'}$. By Kummer theory, the d -root of g defines a finite cyclic covering $\tilde{C}_0$ of C_1 . By the composite $\tilde{C}_0 \rightarrow \mathbf{F}_{q'} \rightarrow \mathbf{F}_q$, $\tilde{C}_0$ is considered as a curve on $\mathbf{F}_q$. Note that it is not always geometrically connected. Let $\pi : \tilde{C}_0 \rightarrow C_0$ be the natural projection. Then we have

$$G = \text{Aut}(\tilde{C}_0/C_0) \simeq \mu_d \rtimes \text{Gal}(\mathbf{F}_{q'}/\mathbf{F}_q).$$

The induced representation $\text{Ind}_{\mu_d}^G(\chi)$ is denoted by Ind and the group ring $\bar{\mathbf{Q}}_l[G]$ is denoted by A . The Kummer sheaf K_0 on C_0 is defined by

$$K_0 = \pi_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind}.$$

Then it is easy to see that

$$K_0|_C \simeq \bigoplus_{\sigma \in \text{Gal}(\mathbf{F}_{q'}/\mathbf{F}_q)} \text{Kum}(g, \chi^\sigma).$$

Let $\tilde{X}_0 = X_0 \times_{C_0} \tilde{C}_0$ and $\tilde{X} = \tilde{X}_0 \otimes_{\mathbf{F}_q} \mathbf{F}$. The projection $\tilde{X}_0 \rightarrow C_0$ and $\tilde{X} \rightarrow C$ is denoted by $\tilde{f}_0$ and $\tilde{f}$ respectively. Then the group G acts on $\tilde{X}_0$ and $\tilde{X}$. Let $W_0 = C_0 - \{s_0\}$, $U_0 = W_0 - \text{Supp}(g)$ and $W = W_0 \otimes_{\mathbf{F}_q} \mathbf{F}$, $U = U_0 \otimes_{\mathbf{F}_q} \mathbf{F}$. The natural inclusions $U \rightarrow W$, $W \rightarrow C$ and $U \rightarrow C$ are denoted by j_1 , j_2 and j_3 respectively.

Proposition 2.3. The action of the Frobenius $\text{Frob}_{\mathbf{F}_q}$ on

$$(2.1) \quad H^1(C, (j_2)_*(j_1)_!(R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind}) \otimes \mathcal{F})$$

is pure of weight $i + 1$.

Proof. Let $\tilde{X}^0 = X_0 \times_{\mathbf{F}_{q'}} \mathbf{F}$ be the connected component of $\tilde{X}$, and $\tilde{f}^0 : \tilde{X}^0 \rightarrow C$ be the natural projection. Then the $\text{Gal}(\mathbf{F}/\mathbf{F}_{q'})$ -module (2.1) is isomorphic to the induced representation of the $\text{Gal}(\mathbf{F}/\mathbf{F}_{q'})$ -module

$$(2.2) \quad H^1(C, (j_2)_*(j_1)_!(R^i \tilde{f}_*^0 \bar{\mathbf{Q}}_l \otimes_{\mathbf{Q}_l[\mu_d]} \bar{\mathbf{Q}}_l(\chi)) \otimes \mathcal{F})$$

Therefore it is enough to prove that (2.2) is pure of weight $i + 1$ under the action of $\text{Gal}(\mathbf{F}/\mathbf{F}_{q'})$. By the condition of Lemma 2.1 (4), we have

$$(j_1)!(R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi) \otimes \mathcal{F}) \simeq (j_1)_*(R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi) \otimes \mathcal{F}).$$

Therefore we have

$$(j_2)_*(j_1)!(R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi) \otimes \mathcal{F}) \simeq (j_3)_*(R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi) \otimes \mathcal{F}).$$

Since f is projective smooth on U , $R^i f_* \bar{\mathbf{Q}}_l \otimes \text{Kum}(g, \chi) \otimes \mathcal{F}$ is punctually pure of weight i . Therefore by the theorem of purity in [D], Theorem 3.2.3, we get the proposition.

Now we fix an identification $\bar{\mathbf{Q}}_l \xrightarrow{\iota} \bar{\mathbf{Q}}_{l'}$ for primes l and l' different from p . Let $D_0 \rightarrow C_0$ be a Galois covering with the finite Galois group G and τ be a $\bar{\mathbf{Q}}_l$ -valued representation of G . Let $\mathcal{F}_0(\tau)$ be the l -adic sheaf associated to the representation τ . The restriction of $\mathcal{F}_0$ to C is denoted by $\mathcal{F}$. We compare elements of $\bar{\mathbf{Q}}_l$ and $\bar{\mathbf{Q}}_{l'}$ via the isomorphism ι .

Proposition 2.4. *The characteristic polynomial of $\text{Frob}_{\mathbf{F}_q}$ on*

$$H_c^1(U, R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}(\tau))$$

is independent of l .

Proof. By Remark 2.2 and Lefschetz trace formula, we have the following equality for zeta function of the sheaf $R^i(\tilde{f}_0)_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}_0(\tau)$ on U_0 .

$$\begin{aligned} & \det(1 - t \text{Frob}_{\mathbf{F}_q} \mid H_c^1(U, R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}(\tau))) \\ &= \prod_{x \in |U_0|} \det(1 - t \text{Frob}_x \mid (R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}(\tau))_{\bar{x}})^{-1} \\ &= \prod_{x \in |U_0|} \det(1 - t \text{Frob}_x \mid H^i((\tilde{f})^{-1}(\bar{x}), \bar{\mathbf{Q}}_l) \otimes_A \text{Ind} \otimes \mathcal{F}(\tau)_{\bar{x}})^{-1} \end{aligned}$$

Since $\tilde{f}$ is projective smooth at $x \in U$, the right hand side is independent of l by the classical Weil conjecture and relation between Zeta function and the Frobenius action and the action of G on the $\mathbf{F}$ rational points of $\tilde{f}^{-1}(x)$. Therefore we get the proposition.

§3 PROOF OF THE MAIN THEOREM

Let K_0 be a local field of characteristic $p > 0$ with a residue field $\mathbf{F}_q$. Let X_η be a projective geometrically smooth variety of dimension n over K_0 . The variety X_η is called globalizable if there exist projective varieties C_0, X_0 over $\mathbf{F}_q$ of dimension one and $n + 1$, a morphism $g : X_0 \rightarrow C_0$, a point $p \in C_0(\mathbf{f}_q)$ and an isomorphism of local field between K_0 and the quotient field K_p of the completion of the structure sheaf of C_0 at p such that the base change of X_0 by the morphism $\text{Spec}(K_0) \rightarrow \text{Spec}(K_p) \rightarrow C_0$ is isomorphic to X_η . Let τ be a representation of $\pi_1(C_0)$ with a finite image and $\bar{\mathbf{Q}}_l(\tau)$ be the representation space of τ .

Proposition 3.1. *The characteristic polynomial of the Frobenius action $\text{Frob}_{\kappa(s_0)}$ on the sapce $(H^i(X_{\bar{\eta}}, \mathbf{Q}_l) \otimes \bar{\mathbf{Q}}_l(\tau))^{\text{Gal}(\bar{K}_p/K_p)}$ is independent of l .*

Proof. To show the independence of l , it is enough to compare two primes l_1 and l_2 . By using Lemma 1.1, we may assume that $X_{\bar{\eta}}$ is globalizable. We use the same notation for X_0, C_0, p_0 and so on. First we choose a rational function g on C_0 , an integer d , and a character $\chi : \mu_d(\mathbf{F}) \rightarrow \mathbf{Q}_l^\times$ satisfying the properties in Lemma 2.1. We consider varieties $\tilde{C}_0, \tilde{C}, \tilde{X}_0, \tilde{X}$ and so on as in §2. Now we consider the exact sequence of sheaves on C :

$$\begin{aligned} 0 \rightarrow (j_3)_!(R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind}) \otimes \mathcal{F}(\tau) &\rightarrow (j_2)_*((j_1)_!(R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind}) \otimes \mathcal{F}(\tau)) \\ &\rightarrow i_s^*((j_3)_*(R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}(\tau))) \rightarrow 0, \end{aligned}$$

where $i_s : \{s\} \rightarrow C$ is the natural inclusion. Taking the cohomology, we have the long exact sequence;

$$\begin{aligned} 0 \rightarrow H^0(\text{Spec}(K), H^i(\tilde{X}_{\bar{\eta}}, \bar{\mathbf{Q}}_l) \otimes_A \text{Ind} \otimes \bar{\mathbf{Q}}_l(\tau)) &\rightarrow H_c^1(U, R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}(\tau)) \\ &\rightarrow H^1(C, (j_2)_*(j_1)_!(R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind} \otimes \mathcal{F}(\tau))) \rightarrow 0. \end{aligned}$$

The weight of

$$(3.1) \quad H^0(\text{Spec}(K), H^i(\tilde{X}_{\bar{\eta}}, \bar{\mathbf{Q}}_l) \otimes_A \text{Ind} \otimes \bar{\mathbf{Q}}_l(\tau))$$

is less than or equal to i , and that of $H^1(C, (j_2)_*((j_1)_!(R^i \tilde{f}_* \bar{\mathbf{Q}}_l \otimes_A \text{Ind}) \otimes \mathcal{F}(\tau)))$ is purely $i + 1$. Therefore the characteristic polynomial of the Frobenius action on (3.1) is independent of l . By the property Lemma 2.1 (2) of the choice of g , (3.1) is isomorphic to

$$H^0(\text{Spec}(K), H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l) \otimes \bar{\mathbf{Q}}_l(\tau))^{\oplus r},$$

where $r = [\mathbf{F}_q(\mu_d) : \mathbf{F}_q]$, and we get the theorem.

By using Poincare duality and local duality, we have the following corollary.

Corollary 3.2. *The characteristic polynomial of the Frobenius action on $H^1(\text{Spec}(K), H^i(X_{\bar{\eta}}, \mathbf{Q}_l) \otimes \bar{\mathbf{Q}}_l(\tau))$ is independent of l .*

Now we recall some properties of the weight monodromy filtration $W_\bullet$. Let G_0 and G be the absolute Galois groups of K_0 and K respectively. There exists an open compact subgroup $\tilde{M}^0$ which acts on $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)$ unipotently and the corresponding quotient of $\tilde{M}^0$ is denoted by M^0 . It is isomorphic to $\mathbf{Z}_l(1)$. Then the image M of G in $\text{Aut } H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l)$ contains M^0 and we may assume that M^0 is a normal subgroup of M by changing M^0 sufficiently small. Then the quotient group M/M^0 acts on $\mathbf{Z}_l(1)$ by the conjugation and the corresponding character is denoted by β . Then the relation of $N : Gr_k^W \rightarrow Gr_{k-2}^W$ and $g \in M/M^0$ is given by $Ng = \beta(g)^{-1}gN$. Therefore, by the universal property of the filtration W , the action of the group M/M^0 preserves the filtration W and as a consequence M/M^0 acts on the associate graded module $Gr_k^W = Gr_k^W(H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))$. Again by the relation of N and g , the group M/M^0 acts on the primitive part P . The corresponding representation of

P_k is denoted by α_k . Then as representations of M/M^0 , Gr_k^W , $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)^{M^0}$, and the coinvariant $H^i(X_{\bar{\eta}}, \mathbf{Q}_l)_{M^0}$ under the action of M^0 are isomorphic to

$$\begin{aligned} Gr_k^W &= \oplus_{m \geq 0} \alpha_{k+2m} \otimes \beta^m, \\ H^i(X_{\bar{\eta}}, \mathbf{Q}_l)^{M^0} &= \oplus_{m \geq 0} \alpha_m \otimes \alpha^m, \\ H^i(X_{\bar{\eta}}, \mathbf{Q}_l)_{M^0} &= \oplus_{m \geq 0} \beta_m. \end{aligned}$$

It is easy to see that all the intersection of the kernel of α and α_k corresponds to the maximal nilpotent subgroup for the action of M on $\text{Aut}(H^1(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))$. Changing notation, this group is denoted by M^0 . Let $\tilde{M}^0$ be the inverse image of M^0 under the natural map $G \rightarrow M$. Let N_0 be an open compact subgroup of G_0 where the intersection $N_0 \cap G$ is contained in $\tilde{M}_0$. The corresponding extension of K_0 is denoted by L_0 . The residue field of L_0 is denoted by $\mathbf{F}_{q'}$.

Theorem 3.3.

- (1) *The representation α_k and β of G is independent of l . Especially, $\tilde{M}^0$ does not depend on l and the dimension of P_k is independent of l .*
- (2) *The characteristic polynomial of the Frobenius action $\text{Frob}_{\mathbf{F}_{q'}}$ on P_k is independent of l .*

Proof of (1). Let τ be a finite dimensional irreducible representation with a finite image H . Then there exists a Galois covering $\pi : D \rightarrow C$ and a lifting q of p such that the quotient field $K_{D,q}$ of the completion of the structure sheaf $\mathcal{O}_D$ of D at q corresponds to the quotient H of G . Then the curve C , D and all the morphism in $\text{Gal}(D/C)$ is defined over a finite extension $\mathbf{F}_{q'}$ of $\mathbf{F}_q$. The model of C and D defined on $\mathbf{F}_{q'}$ is denoted by C_1 and C_2 respectively. Let C'_1 be the covering between $D \rightarrow C$ corresponding to the stabilizer $\text{Stab}_q(G)$ of q . Since $H \simeq \text{Stab}_q(G) \simeq \text{Gal}(D_1/C'_1)$, we can consider a sheaf $\mathcal{F}'_1(\tau)$ on C'_1 . We apply Proposition 3.1 to $\mathcal{F}'_1(\tau)$ and C'_1 . Then the characteristic polynomial of the Frobenius action on $(\bar{\mathbf{Q}}_l(\tau) \otimes H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))^G$ is independent of l . Let $\tilde{M}_l^0$ be the maximal unipotent subgroup for the representation $H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l)$. Then there exists a finite normal subgroup $\tilde{M}_l^1$ of $\tilde{M}_l^0$ such that the restriction of τ to $\tilde{M}_l^1$ is trivial. Then the characteristic polynomial of Frobenius action on

$$\begin{aligned} &(\bar{\mathbf{Q}}_l(\tau) \otimes H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))^G \\ &= ((\bar{\mathbf{Q}}_l(\tau) \otimes H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))^{\tilde{M}_l^1})^{G/\tilde{M}_l^1} \\ &= (\bar{\mathbf{Q}}_l(\tau) \otimes (H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))^{\tilde{M}_l^1})^{G/\tilde{M}_l^1} \\ &= (\bar{\mathbf{Q}}_l(\tau) \otimes (H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l))^{\tilde{M}_l^0})^G \\ &= \text{Hom}_G(\bar{\mathbf{Q}}_l(\tau^*), H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}})^{\tilde{M}_l^0}) \end{aligned}$$

is independent of l . The dimension of weight k -part of the above equality is nothing but the multiplicity of τ^* in $\alpha_k \otimes \beta^k$. Since τ is the arbitrary irreducible representation of G with finite image, $\alpha_k \otimes \beta^k$ is independent of l . Similarly, the characteristic polynomial of Frobenius action on

$$\begin{aligned} &H^1(\text{Spec}(K), H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l) \otimes \bar{\mathbf{Q}}_l(\tau)) \\ &= \text{Hom}_G(\bar{\mathbf{Q}}_l(\tau^*), H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l) \otimes \bar{\mathbf{Q}}_l(\tau)) \end{aligned}$$

is independent of l and we have the l -independence of α_k ($k \geq 0$). This proves (1) of Theorem 3.3. Therefore by the characterization of $\tilde{M}_l^0$ gives as above, it is independent of l .

Let D be a covering of C and $q \in D$ a lifting of p such that the field $K_{D,q}$ defined as above corresponds to the sub group $\tilde{M}^0$ of G . Then D, C are defined over some finite extension $\mathbf{F}_{q'}$ of $\mathbf{F}_q$ and the model over $\mathbf{F}_{q'}$ are denoted by D_1 and C_1 respectively. We may assume $q \in D_1(\mathbf{F}_{q'})$. We apply Proposition 3.1 to $q \in D_1$ and the trivial representation of $\pi_1(D_1)$. Then the characteristic polynomial of Frobenius action on $H^i(X_{\bar{\eta}}, \bar{\mathbf{Q}}_l)^{\tilde{M}^0}$ is independent of l . Considering weight of P_k , the characteristic polynomial of Frobenius on P_k is independent of l .

Remark 3.4.

- (1) *By the independence of l for P_k implies the independence for the Swan conductor.*
- (2) *Since β is independent of l , it is order 2 character by Chebotarev density theorem.*

REFERENCES

- [J] de Jong, A.J., *smoothness, semi-stability and alterations*, Publ. Math. IHES **83** (1996), 51-93.
- [D] Deligne, P., *La conjecture de Weil. II*, Publ. Math. IHES **52** (1981), 313-428.